

Cruise Report
Cruise no. 1410

**Joint investigations on blue whiting south of the Faroes and in the area
west of the British Isles**

27 March - 9 April 2014

R/V Magnus Heinason OW2252



Jan Arge Jacobsen
Leon Smith
Poul Vestergaard
Jens Arni Thomassen



HAVSTOVAN
FAROE MARINE RESEARCH INSTITUTE
POBox 3051 - FO 110 Tórshavn, Faroe Islands

INTRODUCTION

The main aims of this survey was to investigate the distribution and abundance of spawning and post-spawning blue whiting in the areas south of the Faroes, west of the British Isles and Porcupine Bank. Zooplankton and hydrographic data were collected along the cruise tracks.

The cruise was part of the joint international blue whiting survey (IBWSS) on the spawning grounds west of the British Isles, the Porcupine Bank and the Rockall Bank. Four parties and six research vessels (see text table below) took part in the survey, coordinated by the “Working Group of International Pelagic Surveys” (WGIPS) in ICES (formerly WGNAPES). The results from all vessels combined will be used in the assessment of blue whiting by the “Working Group on Widely Distributed Stocks” (WGWIDE) in August 2014.

Ship	Nation
M. Heinason	Faroes
G.O. Sars	Norway
F. Nansen	Russia
Celtic Explorer	Ireland (EU)
Tridens	Netherlands (EU)

The present survey report is based on data from R/V *Magnus Heinason* only. Therefore no estimate of blue whiting is given due to incomplete coverage of the whole spawning area.

MATERIAL AND METHODS

Cruise tracks with hydrographic stations (CTD) and pelagic trawl stations in the surveyed area are shown in **Fig. 1**. Acoustic data were recorded with a Simrad EK-60 echo sounder. Data from the hull mounted 38 kHz transducer were logged at sea and used in the fish abundance estimation. The area backscattering recordings (s_A) per nautical mile were averaged by each nautical mile and the recordings were scrutinised on a daily basis with the EchoView 5 software and allocated to blue whiting, plankton or other fish (e.g. pearlside, lantern fish) based on pelagic trawling aimed at the various acoustic recordings. The 38 kHz Echo sounder was calibrated prior to survey with a standard copper sphere.

RESULTS

The preliminary results from the Faroese investigations in April 2014 indicated large quantities of small (one year old) immature blue whiting in the Faroese area. South of the Wyville-Thomson Ridge and south of the banks larger mature blue whiting was found. The main bulk of the spawning stock of blue whiting was still south of the area surveyed by *Magnus Heinason*, and had by early May not reached the Faroese zone on its northward post-spawning migration towards the Nordic Seas to feed. The sum of the s_A values of blue whiting per each nautical mile along the cruise tracks from the *Magnus Heinason* survey are shown in **Fig. 2**, and the average s_A values of blue whiting by statistical squares in **Fig. 3**. The high abundance of immature 1 and 3 year old blue whiting in the area south of the Faroes is a promising sign for stock.

The length distribution of blue whiting is shown in **Fig. 4**. The mean length was 22 cm (mean weight 82 g). However, there were two tops in the length distribution, one around 18 cm (1 year old) and one around 25 cm (mainly 3 year olds and older fish). The age distribution is shown in **Fig. 5**, and the young (1-3 yr) immature fish was mainly found in the northern area (around the banks and north of the Wyville-Thomson Ridge) while the older fish were found further south. A combined abundance estimates of blue whiting will be calculated at a post-survey meeting later in April 2014 and reported to ICES in September 2014.

The sea-surface temperature (SST) in the surveyed area south of the Faroes was between 7-9°C (**Fig. 6**). Temperature and salinity casts down to 1000 m if possible were taken along the track. The zooplankton samples generally showed very low abundance, indicating that the spring bloom was in its beginning.

Other species

Dealfish (*Trachipterus arcticus*) was only caught on a few stations, while pearlside (*Maurolicus mülleri*) was abundant in the upper scattering layer and lanternfish was abundant in the deeper layers throughout the surveyed area.

Survey effort for *Magnus Heinason* 27/3-9/4 2014:

Effective survey period	Length of cruise track (nm)	Trawl stations	CTD stations	Plankton sampling	Aged fish	Length-measured fish
29/3-8/4	1400	10	21	21	338	1143

Trawl specifications for *Magnus Heinason*:

Circumference (m)	640
Vertical opening (m)	45–55
Mesh size in codend (mm)	40
Typical towing speed (kn)	3.0–4.0




Figure 1. Cruise tracks (black lines) with hydrographic stations (light yellow circles) and trawl stations (green circles) south of the Faroes, *Magnus Heinason* cruise 1410, 27/3-9/4 2014.




Figure 2. Integration values (s_A , m^2/nm^2) of blue whiting per each nm along the cruise tracks, *Magnus Heinason* cruise 1410, 27/3-9/4 2014. The size of the circles corresponds to amount of fish.




Figure 3. Mean integration values (s_A , m^2/nm^2) of blue whiting per statistical square (1x2 degrees), *Magnus Heinason* cruise 1410, 27/3-9/4 2014.




Figure 4. Length distribution of blue whiting south of the Faroes, *Magnus Heinason* cruise 1410, 27/3-9/4 2014.




Figure 5. Age distribution of blue whiting south of the Faroes, *Magnus Heinason* cruise 1410, 27/3-9/4 2014.




Figure 6. Sea-surface temperature (°C) south of the Faroes, *Magnus Heinason* cruise 1410, 27/3-9/4 2014.