



CEnd 10/18 *Nephrops* TV Survey Final Report - 2018

Title: Farn Deepes *Nephrops* Grounds (FU6) 2018 UWTV Survey Report.

Scientific crew: Chris Firmin (SIC)
Rosana Ouréns (2SIC)
Karen Vanstaen
Andy Lawler
Jon Elson
Alessio Veneziano
Dan Clarke
Marc Whybrow (MIST)
Alfie Howat (MIST)

Date/Times: Sailing: 19/06/2018 (01:30 BST) / Docking: 26/06/2018 (22:00 BST)

Keywords: *Nephrops norvegicus*; stock assessment; geostatistics; underwater television (UWTV); benthos

Abstract: This report provides the main results and findings of the 22nd annual underwater television survey on the 'Farn Deepes grounds' ICES assessment area, Functional Unit 6. The survey was multi-disciplinary in nature collecting UWTV video transects and other ecosystem data. The survey design consists of a randomised fixed grid of 110 stations where at each station a sledge mounted TV camera is deployed and a clear 10 minute tow is recorded digitally. In June 2018, 109 stations (TVID) were successfully surveyed in the Farn Deepes area with the TV sledge, from 19 Jun to 26 Jun. In addition to these stations an additional 16 exploratory stations were completed. 1 station had to be dropped due to very close proximity of pipe-laying works and 2 others were relocated slightly from their usual positions 15 stations were repeated due to weather conditions affecting ground contact of the sledge and resulting footage however generally the water clarity was good to excellent throughout the footage recorded.. Burrows were counted by each minute block for 7 clear minutes. The counting performance of the 2017 counters was generally very high, with a Linn's CCC scored average of 0.66. Preliminary results show that the abundance has increased since last year, with abundance now just above the Btrigger threshold as determined by ICES. As with previous years the high abundance area is distributed in the west side of the ground.

Introduction

The Norway lobster (*Nephrops norvegicus* L. 1758) has a wide area of distribution across European waters (from Iceland to the southern coast of Portugal, Morocco and the Mediterranean) and are managed within the scope of the International Council for the Exploration of the Sea (ICES). The *Nephrops* stock assessments are run annually, where catch options are defined for each functional unit (FU) and accordingly on advice from ICES the European Commission sets annual total allowable catches (TAC's) for this species at an ICES sub-area level.

The lack of age-structured data in addition to uncertain historic landings for a number of stocks makes the use of standard stock assessments and forecasting methods, based on commercial catch data, very difficult to apply and unreliable. Additionally, *Nephrops* spend a great deal of time in their burrows and their emergence behaviour is influenced by several factors: time of year, light intensity, tidal strength, etc. So, over the last 22 years in the Farn Deep, assessments for *Nephrops* have become progressively more reliant upon Underwater TV (UWTV) surveys which have enabled the development of fishery independent indicators of stock size, exploitation status and catch advice. This method was firstly implemented in 1992 by Marine Scotland on the Fladen ground, and has subsequently been put into practice by other countries such as Ireland, England, Denmark and Sweden. The UWTV surveys are now listed regularly in 15 ICES Functional Units, being widely used in the North Sea.

The standard methodology involves the use of a sledge mounted camera to film the seabed at a grid of stations conducting TV tows for 10 minutes. Each country has adopted different sampling designs, from random stratifications of the stations up to fix grids, which better fits the grounds. The aim is to identify and count the number of *Nephrops* burrow systems falling within a fixed field of view, along transects of known length. Counts of burrow systems are converted into densities at each station using the width of view and the length of the tow. Each system is assumed to represent one adult *Nephrops* with occupancy assumed to be 100%. Overall abundance is then estimated by raising the mean density to the appropriate strata area or by using geostatistical methods, and total survey abundance, variance and confidence limits are then calculated.

In deep waters the UWTV surveys are still not being used as a standard assessment procedure, due to the complexity of running a sledge at those depths. Alternatives to sledges have been experimented for example by IPMA in Portugal by fitting a camera to the trawl cable; disadvantages of this method are the speed of recording and the angle of the camera, making the visual identification of burrow systems very challenging.

Although this assessment method has been improved over the years there are still some constraints associated with this method. Misidentification of *Nephrops* burrows, high density of burrows, edge effects, clarity and variability of the counters are some of the sources of bias that have been identified in the past and addressed in specialized ICES study groups for *Nephrops* TV surveys. Progress was made in 2009 ICES Benchmark where the main sources of bias were estimated for each functional unit and an overall bias correction factor introduced adjusting the estimates of abundance.

The present survey focuses in the North Sea at the Farn Deep (FU6) area, in the NE coast of England (Figure 1). Total landings for the 2017/2018 season for this area were 1337 tonnes.

Cefas has performed annual UWTV surveys in the Farn Deep area since 1996 (Table 1).

Table 1 – Summary of the UWTV results, since 1997, showing number of valid stations, mean density per meter square, abundance, confidence interval and the method used to estimate the abundance.

Year	Stations	Season	Mean density (burrows/m ²)	Absolute Abundance (millions)	95%CI (millions)	Method
1997	87	Autumn	0.46	1500	125	Box
1998	91	Autumn	0.33	1090	89	Box
1999	-	Autumn	No survey	Box		
2000	-	Autumn	No survey	Box		
2001	180	Autumn	0.56	1685	67	Box
2002	37	Autumn	0.33	1048	112	Box
2003	73	Autumn	0.33	1085	90	Box
2004	76	Autumn	0.43	1377	101	Box
2005	105	Autumn	0.49	1657	148	Box
2006	105	Autumn*	0.37	1244	114	Box
2007	105	Autumn*	0.28	858	23	Geostatistics
2008	95	Autumn*	0.31	987	39	Geostatistics
2009	76	Autumn*	0.22	682	38	Geostatistics
2010	95	Autumn*	0.25	785	21	Geostatistics
2011	97	Autumn*	0.28	878	17	Geostatistics
2012	97	Autumn*	0.24	758	13	Geostatistics
2013	110	Summer	0.23	706	18	Geostatistics
2014	110	Summer	0.24	755	18	Geostatistics
2015	110	Summer	0.18	568	13	Geostatistics
2016	110	Summer	0.24	697	19	Geostatistics
2017	110	Summer	0.29	909	21	Geostatistics
2018	109	Summer	0.30*	954*	tbc	Geostatistics

* *Provisional result*

The specific objectives of the 2018 survey are listed below:

1. To conduct a standard underwater TV survey of Nephrops burrow densities on the Farn Deep grounds, 55 ° 35' - 54 ° 45' N and 1 ° 30' - 0 ° 40' W, (shown in black on figure 1) and to evaluate Nephrops abundance (110 stations).
2. Secondary: To conduct a standard underwater TV survey of additional stations on the Botney Gut / Silver Pit grounds 53 ° 45' - 54 ° 15' N and 1 ° 25' - 2 ° 55' E (shown in figure 2) if time allows.
3. Secondary: To conduct a standard underwater TV survey of additional stations on the Farn Deep grounds (as per objective 1, highlighted red on figure 1), if time allows. These stations are surveyed by NEIFCA as part of their annual autumn TV survey of inshore grounds. By surveying the same stations during the early summer we will be able to estimate seasonal variations in burrow densities.




Figure 1 – Map showing the location of the surveyed area in the Function Unit 6 area (110 stations) and the 15 additional NEIFCA stations.




Figure 2 - C End 10/18 final stations for Silver Pit / Botney Gut area (FU5) – to west of greyed area.

1. Material and Methods

The 2018 North Sea *Nephrops* UWTV survey took place on RV Endeavour between 19th to 26th June. The departure and arrival port was Lowestoft.

TV survey – Survey design

For the Farn Deep the survey design is based on a randomised fixed grid and includes a total of 110 stations. The initial ground perimeter has been delimited by the combination of VMS data and BGS sediment maps.

At each station a sledge mounted TV camera was deployed and a clear 10 minute tow was recorded to MP4 video files, recorded directly to two separate drives to provide a backup. Vessel position (DGPS) and position of sledge (using a USBL transponder) were recorded every 10 seconds.

The camera system, lights, lasers, altimeter and compass used on the sledge were all manufactured by STR. The camera, lights and lasers were controlled by an onboard mux box, which communicated with a MUX box on board. Camera cable B (coax) was used to receive the live footage from the video camera. This is the second year using this camera system, offering better video quality than the SIMRAD camera used in previous years, although there were the occasional issues with jerky footage.

The sledge was equipped with (Figure 3):

- An IP camera (720p) at an oblique angle to the sea bed, sighted towards the front of the sled.
- The sledge was mounted with 6 LED lights: 2 LED lights on either side plus 2 LED lights at the front to fully illuminate the field of view.
- Two green fan lasers to delimit the field of view (field of view 81.5 cm);
- A transponder so that the sledge can be retrieved if lost;
- The ESM2 logger, to record turbidity readings, depth; salinity and oxygen levels;
- With the new system, an independent altimeter and compass are used, which log to a txt file.




Figure 3 – Sledge used during CEnd 10/18, showing the equipment setup. *Photos by Chris Firmin (Cefas).*

The Dynamic Positioning system (DP) was used throughout the survey to provide a controlled towing speed of around 0.7 knots.

Recounts

In line with SGNEPS recommendations all scientists were trained/re-familiarised using training material and validated using reference footage (measured by Linn's concordance correlation coefficient (CCC)) prior to recounting June 2018 footage. A limit of 0.5 was used to identify counters who need further training. On completion of this process, all CEnd 10/18 recounts were conducted, as blind counts, by two persons during the survey. Here, the number of *Nephrops* burrow systems and the activity in and out of the burrows were counted by each minute block (for 7 clear minutes). In case the field of view became obscured by cloud the seconds obscured were recorded and all minute blocks with more than 20 minutes obscured were rejected. After all counts were completed, Linn's CCC (with a threshold of 0.5) was again applied to check which stations needed to be reviewed and where a 3rd counter needed to be added.

Whilst reviewing the videos, the visibility, ground type, trawl marks, occurrence of bio-fauna, ground contact of the sledge, cloud and any other interference was recorded during each one-minute intervals, using a classification key.

For analysis, counts of burrow systems are converted into densities at each station using the width of view (81.5 cm) and the length of the tow (extracted from tower position vessel logging). Each system is assumed to

represent one adult *Nephrops* and occupancy is assumed to be 100%. To estimate the spatial structure of *Nephrops* densities a geo-statistical analysis is carried out in the whole area and the total survey abundance, variance and confidence limits are then calculated.

Health and Safety

As required all staff had a valid ENG1 health certificate, Personal Sea Survival Certificate and a valid 'working near water' safety course.

Also the following risk assessments were acknowledged:

- ✓ RA3022 *Nephrops* TV Cruise
- ✓ RA4038 (HS26 MPM-MAS-MIS-RA-16) Cameras with Laser Scaling Systems
- ✓ RA003 Access to and Work on RV Cefas Endeavour
- ✓ RA006 Deploying and Recovering Scientific Equipment (grabs, cameras, buoys) on vessels
- ✓ RA4052 (HS26 MPM-MAS-MOS-RA-05) Unloading and loading vehicles at ports docksides
- ✓ FD-C&F-SHELL-SOP-01 MB001 NEPTVBurrowCount SOP V1.5
- ✓ CP001-7 Fieldwork 'Including working in, on or near to water'
- ✓ CP001-17 Driving for Cefas
- ✓ FD-C&F-SHELL-SOP-01 MB001 NEPTVBurrowCount SOP V1.5.DOC (updated during the survey).
- ✓ G02 – Travelling while on official duty in Official or private vehicles, including loading and unloading equipment, baggage, etc, but excluding the carriage of dangerous chemicals, the use of HGV or specialised vehicles;
- ✓ G03 – Participation in research cruises on CEFAS owned and managed ships. The collection of samples and data all subsequent processing whilst on-board, including the use of the ships sea-rider.
- ✓ FD-CF-SHELL-RA-09-MB001 – *Nephrops* TV cruise activities

Technical aspects/failure

- An STR High Definition 1080p camera systems using fibre-optic umbilical cable was proposed for use during this year's the survey, however the cable had to be repaired and re-terminated after damage sustained during C End 0918. Sustained effort on the part of the MIST engineers ensured this was carried out during the 24 hour changeover, however further testing once at sea revealed a defective MUX termination requiring and replacement parts. Attempts to set up this system were therefore abandoned in favour of the previously-used IP 720p camera with coax cable
- Windows 10 Enterprise machines provided very limited flexibility for data-sharing during the trip. Attempts to mount the survey database on the ship's network were also abandoned due to network issues. Local copies of the database were used with transfer of files around by external hard disc drives.

2. Results and Final Considerations

In June 2018, 109 of the 110 stations (TVID) of the standard survey grid, as well as an additional 16 stations of the NEIFCA survey grid, were successfully surveyed in the Farn Deep (FU6) with the TV sledge from 19th Jun (01:30 BST) to 26th June (22:00 BST). No time was lost due to weather conditions; however high swells over a 16 hour period between 20 June and 21 June resulted in poor footage on 15 stations that had to be repeated. FThe

visibility was, as with the previous two years, generally good, 97% of footage was classed as “Good”, 3% as “Moderate”, and less than 0% as “Poor” (table 1). This time of the year once again proved to be ideal to do the survey as all conditions are more favourable to run a more efficient survey and it allowed also to save time as much fewer stations needed to be revisited due to bad weather conditions and/or poor water clarity.

Table 1 - Classification of UWTV Survey footage 2011–2017.

	Good	Moderate	Poor/ none
2011 – Autumn	49%	48%	2%
2012 – Autumn	74%	22%	4%
2013	95%	5%	0%
2014	91%	9%	0%
2015	99%	1%	0%
2016	92%	8%	0%
2017	93%	7%	0%
2018	97%	3%	0%

Primary objective - TV survey

Nephrops burrow live-counts were made over a 10-minute tow, which was recorded on mp4 files and backed up on two external hard drives. All recordings were then recounted under controlled conditions and the CCC code was used to validate stations and to identify which stations required a 3rd counter, for this propose a threshold of 0.5 was used.

- Burrows were counted by each minute block for 7 clear minutes. The counting performance of the 2018 counters was generally high, with a Linn’s CCC scored average of 0.66.
- As previous years the high abundance area is distributed in the west side of the ground (Figure 4).
- Abundance has now increased from its lowest observed point in the time series in 2016 of 697 million, to 909 million last year and to 954 million this year. The advice will be revised again in the autumn given that the change is more than 1 standard deviation (Table 3).
- Abundance is now slightly above the 2007 trigger level (Figure 5).




Figure 4 – Geostatistical outputs 2009 – 2018, maps of *Nephrops* density distribution (m²)




Figure 5 – *Nephrops* abundance estimates from the UWTV Survey 2001–2017.

Table 2 – Results using the geostatistical model from UWTV-FU 6 *Nephrops* survey in 2007–2017.

Year	Stations	Mean density (burrows/m ²)	Absolute Abundance (millions)	95% confidence interval (millions)
2007	105	0.28	858	23
2008	95	0.31	987	39
2009	76	0.22	682	38
2010	95	0.25	785	21
2011	97	0.28	878	17
2012	97	0.24	758	13
2013	110	0.23	706	18
2014	110	0.24	755	18
2015	110	0.18	568	13
2016	110	0.24	697	19
2017	110	0.29	909	21

The primary objective was over 99% achieved as all but one of the TVID stations successfully surveyed with the TV sledge, all data was inputted and quality checked while onboard and additionally preliminary analysis was made to calculate the abundance estimation for the ground.

The secondary objective of surveying the FU5 Botney Gut / Great Silver grounds was abandoned. Power analysis conducted before the survey had revealed a minimum of 40 stations would have been required for a meaningful result with time available to complete at most 15 after completing the primary objective. Instead, 16 of the 20 NEIFCA stations were surveyed. Two more were abandoned due to the substrate type being unsuitable for the camera sledge and the remaining two were abandoned due to proximity to a cable. These data will be compared with the footage collected by NEIFCA during their annual TV survey which is due to be completed in Autumn 2018.

Final considerations

The main objective of the survey (*Nephrops* abundance estimation) was successfully met for this year in the Farn Deepes. The UWTV coverage was excellent (99% stations done with the TV sledge) and the overall footage quality was very good to excellent in the Farn Deepes grounds due to favourable weather conditions and minimal technical difficulties.

Acknowledgements

We would like to express our thanks and gratitude to the Captain and crew of RV Endeavour for their good will and professionalism during the survey. Also thanks to P&O Maritime for handling all gear and sort any technical difficulties. Finally, thanks to all Cefas staffs onboard for their hard work and enthusiasm in making this survey a success.

Chris Firmin (SIC)